

Supplementary Table 2. Input and output (L/day) between survivors and non-survivors during the early 7 days of ECMO

Variable	Day 1	Day 2	Day 3	Day 4	Day 5	Day 6	Day 7
Survivor							
Input (L)	5.4 (3.7–6.0)	4.1 (3.7–4.5)	4.1 (3.5–5.1)	3.9 (3.5–4.6)	3.8 (3.4–4.7)	4.5 (3.7–4.6)	3.8 (3.5–4.4)
Output (L)	2.1 (1.6–3.2)	2.7 (0.9–4.4)	4.2 (2.3–4.9)	4.9 (4.0–6.9)	4.9 (3.9–5.6)	4.9 (3.6–6.3)	3.9 (3.4–5.0)
Sum (L)	2.1 (1.2–4.7)	1.9 (0.2–2.8)	1.6 (-1.3–2.1)	-1.2 (-1.9–0.1)	-0.8 (-2.1–0.3)	-1.1 (-1.7–0.3)	0.2 (-1.1–0.4)
Non-survivor							
Input (L)	4.3 (3.7–5.3)	5.0 (4.3–5.3)	5.1 (3.9–5.4)	4.5 (4.1–5.6)	3.9 (3.4–5.9)	4.2 (3.6–4.6)	4.1 (3.3–4.4)
Output (L)	1.8 (1.6–2.9)	2.3 (1.4–3.7)	3.9 (2.8–5.3)	4.5 (3.9–5.9)	4.1 (3.2–5.6)	4.8 (3.9–5.6)	3.5 (3.3–5.6)
Sum (L)	2.3 (0.9–2.7)	2.6 (0.7–3.5)	0.2 (-0.3–1.9)	-0.2 (-1.3–0.4)	-0.8 (-0.9–1.9)	-1.0 (-1.6–0.2)	-0.3 (-1.9–0.9)
P-value	0.71	0.37	0.941	0.261	0.112	0.824	0.766

Values are presented as median (interquartile range). During the total seven period, net fluid balance was 2.6 L/day (-3.3 to 3.8 L/day) vs. 10.5 L/day (1.6 to 23.5 L/day) in survivors vs. non-survivors (P=0.053).

ECMO: extracorporeal membrane oxygenation.